

## NOTE

### A Generalization of Fisher's Inequality

Hunter S. Snevily\*

*Department of Mathematics, University of Idaho, Moscow, Idaho 83844*

*Communicated by the Managing Editors*

Received August 26, 1996

In this paper we are concerned with the following conjecture. *Conjecture.* Let  $\mathcal{L}$  be a set of  $k$  consecutive positive integers. In particular, we show this conjecture is true when  $\mathcal{L}$  consists of  $k$  consecutive positive integers. This generalizes a well-known inequality of Fisher's. Our proof simplifies and extends a recent result of Ramanan's. © 1999 Academic Press

[View metadata, citation and similar papers at core.ac.uk](http://core.ac.uk)

## 1. INTRODUCTION

Throughout this paper  $X$  will denote the set  $[n] = \{1, 2, \dots, n\}$  and  $\mathcal{L} = \{l_1, l_2, \dots, l_k\}$  will denote a set of  $k$  arbitrary nonnegative integers. Let  $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$  denote a collection of subsets of  $X$  such that  $|A_i \cap A_j| \in \mathcal{L}$ . We will sometimes refer to  $\mathcal{A}$  as an  $\mathcal{L}$ -intersecting family. In this paper we are, concerned with finding upper bounds for  $|\mathcal{A}|$ . Listed below (in chronological order) are, results related to this problem.

**THEOREM 1** [6]. *In a BIBD (balanced incomplete block design) the number of blocks is never less than the number of points.*

**THEOREM 2** [3]. *If every pair of sets in a uniform family has equal intersection size, then the number of sets does not exceed the number of points.*

**THEOREM 3** [4]. *If  $\mathcal{L} = \{1\}$ , then  $|\mathcal{A}| \leq n$ .*

**THEOREM 4** [11]. *If  $\mathcal{L} = \{\lambda\}$  (for  $1 \leq \lambda \leq n$ ), then  $|\mathcal{A}| \leq n$ .*

\* E-mail: [snevily@uidaho.edu](mailto:snevily@uidaho.edu).

THEOREM 5 [9]. For any  $\mathcal{L}$ ,  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n}{i}$ .

THEOREM 6 [14]. If  $\mathcal{L} = \{1, 2, \dots, k\}$ , then  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n-1}{i}$ .

It can be shown that Bose's result is equivalent to Fisher's result. Thus Theorems 1, 2, and 3 are special cases of Theorem 4 and Theorem 5 was a conjecture of Frankl's and Füredi's. In fact all of the above results are special cases of the following conjecture.

Conjecture 7 [16]. For any set  $\mathcal{L}$  of  $k$  distinct positive integers  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n-1}{i}$ .

To see that Conjecture 7 implies Theorem 5, suppose that Conjecture 7 is true but Theorem 5 is false, i.e. suppose there exists a set  $\mathcal{L}^0 = \{0 = l_1, l_2, \dots, l_k\}$  and a set  $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$  such that  $|A_i \cap A_j| \in \mathcal{L}^0$ , for all  $A_i, A_j \in \mathcal{A}$  but  $|\mathcal{A}| > \sum_{i=0}^k \binom{n}{i}$ . Create a new collection of sets  $\mathcal{A}' = \{A'_1, A'_2, \dots, A'_m\}$  with  $A'_i = A_i \cup \{n+1\}$ . Then  $\mathcal{A}'$  is a collection of subsets of  $X' = X \cup \{n+1\}$  such that  $|\mathcal{A}'_i \cap \mathcal{A}'_j| \in \mathcal{L}^+ = \{1, l_2+1, \dots, l_k+1\}$ , for all  $A'_i, A'_j \in \mathcal{A}'$  and  $|\mathcal{A}'| > \sum_{i=0}^k \binom{(n+1)-1}{i}$ , which is a contradiction.

Although we have not proven Conjecture 7, we have managed to generalize Theorems 1 through 6, as Theorem 9 (given below) will show.

## 2. MAIN RESULT

Let  $\mathcal{L} = \{l_1, l_2, \dots, l_k\}$ , where we shall assume throughout that  $l_1 < l_2 < \dots < l_k$ . Let  $g_{\mathcal{L}}(x) = \prod_{1 \leq i \leq k} (x - l_i)$ , so that  $g_{\mathcal{L}}(x) = 0$  if and only if  $x \in \mathcal{L}$ . Since  $g_{\mathcal{L}}(x)$  is a monic polynomial in  $x$  of degree  $k$ , we can write it (by a change of basis) in the form  $g_{\mathcal{L}}(x) = \sum_{h=0}^k c_h \binom{x}{h}$ , where  $c_0, c_1, \dots, c_k$  are real numbers independent of  $x$ , which we call the *coefficients* of  $\mathcal{L}$ . Let  $C^{\geq 0} = \{h: c_h \geq 0\}$  and  $C^- = \{h: c_h < 0\}$ . Clearly  $k \in C^{\geq 0}$ , since  $c_k = k! > 0$ .

Now suppose  $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$  is an  $\mathcal{L}$ -intersecting family of subsets of  $X = [n]$ . With each  $A_j \in \mathcal{A}$  we associate a real variable  $z_j$ . If  $z = (z_1, z_2, \dots, z_m)$  and  $Y \subset X$  let  $L_Y(z) = \sum_{j: Y \subset A_j} z_j$ . Let  $\binom{X}{h}$  be the set of all  $h$ -element subsets of  $X$ . We shall need the following lemma.

LEMMA 8. If  $z = (z_1, z_2, \dots, z_m)$  and  $y = (y_1, y_2, \dots, y_m)$ , then

$$\sum_{h=0}^k c_h \sum_{Y \in \binom{X}{h}} L_Y(z) L_Y(y) = \sum_{j=1}^m g_{\mathcal{L}}(|A_j|) z_j y_j. \quad (1)$$

*Proof.* It is easy to see that

$$L_Y(z) L_Y(y) = \sum_{\substack{j \\ Y \subseteq A_j}} z_j y_j + \sum_{\substack{i \neq j \\ Y \subseteq A_i \cap A_j}} z_i y_j,$$

so that the LHS of (1) is equal to

$$\begin{aligned} & \sum_{j=1}^m \sum_{h=0}^k c_h \binom{|A_j|}{h} z_j y_j + \sum_{\substack{i, j=1 \\ i \neq j}}^m \sum_{h=0}^k c_j \binom{|A_i \cap A_j|}{h} z_i y_j \\ &= \sum_{j=1}^m g_{\mathcal{L}}(|A_j|) z_j y_j + \sum_{\substack{i, j=1 \\ i \neq j}}^m g_{\mathcal{L}}(|A_i \cap A_j|) z_i z_j, \end{aligned}$$

which is equal to the RHS of (1) because  $|A_i \cap A_j| \in \mathcal{L}$  and  $g_{\mathcal{L}}(x) = 0$  if  $x \in \mathcal{L}$ . ■

We shall be concerned with the following two systems of linear equations:

$$L_Y(x) = 0 \quad \text{for all } Y \subseteq X \quad \text{with } 0 \leq |Y| \leq k \quad (S_1)$$

and

$$L_Y(z) = 0 \quad \text{for all } Y \subset X \quad \text{with } |Y| \in C^{\geq 0}. \quad (S_2)$$

We are now ready to state and prove our main result.

**THEOREM 9.** *If  $\mathcal{A}$  is an  $\mathcal{L}$ -intersecting family of subsets of  $X$ , then*

- (a)  $S_1$  has only the trivial solution (hence  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n}{i}$ ), and
- (b) if  $g_{\mathcal{L}}(|A_j|) \geq 0$  for all  $j$ , then  $|\mathcal{A}| \leq \sum_{i \in C^{\geq 0}} \binom{n}{i}$ .

*Proof.* (a) Suppose that  $\bar{z} = (\bar{z}_1, \bar{z}_2, \dots, \bar{z}_m)$  is solution to  $S_1$ . Set  $z_j = \bar{z}_j$  for all  $j$  and  $y_j = \bar{z}_j$  or  $-\bar{z}_j$  according as  $g_{\mathcal{L}}(|A_j|) \geq 0$  or  $< 0$ . Then the LHS of (1) is 0, and the RHS is  $\sum_{j=1}^m |g_{\mathcal{L}}(|A_j|)| \bar{z}_j^2$ . We deduce immediately that  $\bar{z}_j = 0$  for all  $j$  with  $g_{\mathcal{L}}(|A_j|) \neq 0$ . In particular, this holds whenever  $|A_j| > l_k$ . Let  $\mathcal{L}_i = \{l_1, \dots, l_i\}$  for  $i = 1, \dots, k$ . Since  $\bar{z}_j = 0$  whenever  $|A_j| > l_k$ , the value of  $L_Y(\bar{z})$  is unchanged if we delete all such sets  $A_j$  from the family  $\mathcal{A}$  thereby obtaining an  $\mathcal{L}_{k-1}$  intersecting family. Repeating the above argument with this family, noting that  $\bar{z}$  is evidently a solution to the subset of equations in  $S_1$  corresponding to  $0 \leq |Y| \leq k-1$ , we find now that  $\bar{z}_j = 0$  whenever  $|A_j| > l_{k-1}$ . Continuing in this way with  $\mathcal{L}_{k-2}, \dots, \mathcal{L}_1$ , we find eventually that  $\bar{z}_j = 0$  for all  $j$  with  $|A_j| > l_1$ . But there can be at

most one set in  $\mathcal{A}$  of cardinality  $l_1$ , and since  $L_{\emptyset}(\bar{z}) = 0$  we conclude that  $\bar{z}_j = 0$  for all  $j$ . This completes the proof of (a).

(b) Suppose now that  $\bar{z}$  is a solution to  $S_2$ . Then setting  $z_j = y_j = \bar{z}_j$  for each  $j$ , (1) becomes

$$\sum_{h \in C^-} c_y \sum_{Y \in \binom{C_h}{h}} L_Y(\bar{z})^2 = \sum_{j=1}^m g_{\mathcal{L}}(|A_j|) \bar{z}_j^2. \quad (2)$$

Since the LHS of (2) is nonpositive and the RHS is nonnegative, both sides are 0, and so  $\bar{z}$  is a solution to  $S_1$ . By (a) we deduce that  $S_2$  has only the trivial solution, and (b) follows. ■

We call  $\mathcal{L}$  a *gte-set* (greater-than or-equal to set) if  $g_{\mathcal{L}}(x) \geq 0$  for  $x = l_1, l_1 + 1, \dots, l_k$ . We shall say that the coefficients of  $\mathcal{L}$  *alternate in sign* if  $h$  is alternately in  $C^{\geq 0}$  and in  $C^-$  for  $h = k, k-1, \dots, 0$ .

**COROLLARY 10.** *Suppose that  $\mathcal{A}$  is an  $\mathcal{L}$ -intersecting family and the coefficients of  $\mathcal{L}$  alternate in sign. If  $g_{\mathcal{L}}(|A_j|) \geq 0$  for all  $j$  (which must happen if  $\mathcal{L}$  is a gte-set), then  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n-1}{i}$ .*

*Proof.* Clearly  $|A_j| \geq l_1$  for all  $j$ , and  $g_{\mathcal{L}}(|A_j|) > 0$  if  $|A_j| > l_k$ , and so  $g_{\mathcal{L}}(|A_j|) \geq 0$  for all  $j$  if  $\mathcal{L}$  is a gte-set. Since  $c_k \in C^{\geq 0}$  if the coefficients of  $\mathcal{L}$  alternate in sign then (b) gives

$$|\mathcal{A}| \leq \sum_{i=0}^{\lfloor 1/2k \rfloor} \binom{n}{k-2i} = \sum_{i=0}^n \binom{n-1}{i}$$

by a well-known recurrence for binomial coefficients. ■

**COROLLARY 11.** *Let  $\mathcal{L}$  be a set of  $k$  consecutive positive integers, and suppose that  $\mathcal{A}$  is an  $\mathcal{L}$ -intersecting family. Then  $|\mathcal{A}| \leq \sum_{i=0}^k \binom{n-1}{i}$ .*

*Proof.* Clearly  $\mathcal{L}$  is a gte-set, and so it remains to show that the coefficients of  $\mathcal{L}$  alternate in sign. Suppose  $\mathcal{L} = \{m, m+1, \dots, m+k-1\}$  where  $m \geq 1$ . Then  $g_{\mathcal{L}}(x) = k! \binom{x-m}{k}$ , and so the result follows from the identity

$$\binom{x-m}{k} = \sum_{i=0}^k (-1)^{k-i} \binom{m-1+k-i}{k-i} \binom{x}{i}. \quad (3)$$

To prove (3), choose  $x$  to be an integer greater than  $m+k$  and equate the coefficients of  $y^k$  on the two sides of the equation

$$(1+y)^{x-m} = (1+y)^{-m} (1+y)^x.$$

Since (3) holds as an equation for all such  $x$ , it must be an identity. ■

## 3. CONCLUSION

We should mention in [16] it was shown that, given any set  $\mathcal{L}$  of  $k$  positive integers that Conjecture 7 is true when  $n$  is sufficiently large ( $|X| = n$ ). We offer one more conjecture concerning the extremal families of Conjecture 7.

*Conjecture 12.* If  $\mathcal{L}$  is a set of  $k$  consecutive positive integers and  $\mathcal{A}$  is an  $\mathcal{L}$  intersecting family with  $|\mathcal{A}| = \sum_{i=0}^k \binom{n-1}{i}$ , then  $|\{A: A \in \mathcal{A}\}| \leq k + 1$ .

## ACKNOWLEDGMENTS

This paper was strongly influenced by [1], [12], and [14]. I also thank Rick Wilson for his helpful comments and André Kézdy for his help in preparing this paper. After showing Rick Wilson Theorem 9 (or a variant thereof), he responded with an elegant proof of part (b) using incidence matrices. Finally, I thank Douglas Woodall for providing a much needed clarification of an earlier version of this paper.

## REFERENCES

1. N. Alon, L. Babai, and H. Suzuki, Multilinear polynomials and Frankl-Ray-Chaudhuri-Wilson type theorems, *J. Combin. Theory Ser. A* **58** (1991), 65–180.
2. L. Babai and P. Frankl, “Linear Algebra Methods in Combinatorics I, (preliminary version 2),” Dept. of Comp. Science, Univ. of Chicago, 1992.
3. R. C. Bose, A note on Fisher’s inequality for balanced incomplete block designs, *Ann. Math. Stat.* **20** (1949), 619–620.
4. N. G. deBruijn and P. Erdős, On a combinatorial problem, *Indag. Math.* **10** (1948), 421–423.
5. M. Deza and P. Frankl, Erdős-Ko-Rado theorem—22 years later, *SIAM J. Algebraic Discrete Methods* **4**, No. 4 (1983), 419–431.
6. R. A. Fisher, An examination of the different possible solutions of a problem in incomplete blocks, *Annals Eugenics* **10** (1940), 52–75.
7. P. Frankl, The shifting technique in extremal set theory, in “Proc. of the British Combinatorial Conf., London Math. Soc. Lecture Notes” (C. Whitehead, Ed.), pp. 81–110, 1987.
8. P. Frankl and Z. Füredi, Families of finite sets with missing intersections, *Colloquia Math. Soc. János Bolyai* **37** (1981), 305–320.
9. P. Frankl and R. M. Wilson, Intersection theorems with geometric consequences, *Combinatorica* **1**, No 4 (1981), 357–368.
10. R. L. Graham and H. O. Pollak, On embedding graphs in squashed cubes, in “Graph Theory and Appl., Proc. Conf. Western Mich. Univ., Kalamazoo,” Springer Lecture Notes in Math., Vol. 303, pp. 99–110, 1972.
11. K. N. Majumdar, On some theorems in combinatorics relating to incomplete block designs, *Ann. Math. Stat.* **24** (1953), 377–389.
12. R. Palisse, A short proof of Fisher’s inequality, *Discrete Math.* **111** (1993), 421–422.
13. L. Pyber, An extension of a Frank-Füredi theorem, *Discrete Math.* **52** (1984), 253–268.
14. G. V. Ramanan, Proof of a conjecture of Frankl-Füredi, preprint.

15. H. J. Ryser, An extension of a theorem of deBruijn and Erdős on combinatorial designs, *J. Algebra* **10** (1968), 246–261.
16. H. S. Snevily, On generalizations of the deBruijn–Erdős theorem, *J. Combin. Theory Ser. A* **68** (1994), 232–238.
17. H. Tverberg, On the decomposition of  $K_n$  into complete bipartite graphs, *J. Graph Theory* **6** (1982), 493–494.